

CLOSED-CONSTRUCTIBLE FUNCTIONS ARE PIECE-WISE CLOSED

ALEXEY OSTROVSKY

ABSTRACT. A subset $B \subset Y$ is constructible if it is an element of the smallest family that contains all open sets and is stable under finite intersections and complements. A function $f : X \rightarrow Y$ is said to be piece-wise closed if X can be written as a countable union of closed sets Z_n such that f is closed on every Z_n . We prove that if a continuous function f takes each closed set into a constructible subset of Y , then f is piece-wise closed.

All spaces in this paper are supposed to be separable and metrizable, and all the functions are supposed to be continuous and onto.

A subset $B \subset Y$ is *constructible* if it is an element of the smallest family that contains all open sets and is stable under finite intersections and complements.

In topology, a constructible set is a finite union of locally closed sets (a set is *locally closed* or is an LC_1 -set if it is the intersection of an open set and a closed set).

In algebraic geometry, a constructible set is any zero set of a system of polynomial equations and inequations. A function f is said to be *closed-constructible* (resp., *closed- F_σ*) if f takes closed sets into constructible (resp., F_σ) ones. It is clear that every closed-constructible function is closed- F_σ .

A function $f : X \rightarrow Y$ is said to be *piece-wise closed* if X can be written as a countable union of closed sets Z_n such that f is closed on every Z_n .

Hansell, Rogers and Jayne gave a corrected form of their previous result [6, Theorem 1], [7] using additional hypotheses (a) – (d) [4, Theorem 3].

Under hypothesis (a) (=Fleissner's axiom, which is consistent with the usual axiom ZFC), they established the correctness of their first conclusion [6][Lemma 2]: each continuous, closed- F_σ function between absolute Souslin sets X and Y is piece-wise closed.

Unfortunately, the theory above is not sufficient for important applications such as the case of closed- LC_1 functions between non-Souslin subsets of the real line \mathbf{R} .

Motivated by this observation, we study the extensions of the theory of closed-Borel functions, whose major case is closed- LC_2 functions (a simple case for such non-continuous functions was recently considered in [10]).

We obtain the following main theorem:

Theorem 1. *Every closed-constructible function is piece-wise closed.*

Theorem 1 immediately implies the following:

Corollary 1. *Each constructible-measurable, closed (or open), one-to-one function is piece-wise continuous.*

Note that this result is different from that of Banach and Bokalo [2], which states that a constructible-measurable function of hereditary Baire space X is piece-wise continuous.

Finally, we give some simple observations to the hypothesis (d) mentioned in the beginning: each preimage $f^{-1}(y)$ of points from Y is compact.

This hypothesis looks fairly strong: as we demonstrated in simple Proposition 3, a function f under hypothesis (d) becomes a closed- F_σ function. However, in the same situation (Example 3), such a conclusion becomes false if the requirement (d) is weakened to the following: each preimage $f^{-1}(y)$ of points from Y is completely metrizable.

Proposition 1. *Let $f : X \rightarrow Y$ be a continuous function with compact fibers and f take elements of a clopen base \mathcal{B} into closed sets. Then f is a closed and, hence, closed-constructible function.*

Note that in following Example 1 all preimages $f^{-1}(y)$ of points from Y are completely metrizable, but not complete under the given metric.

Example 1. *There exists a continuous and open function $f : I_X \rightarrow I_Y$ from Polish spaces $I_X, I_Y \subset \mathcal{C}$ that takes elements of some clopen base of I_X into clopen sets, but f is not closed- F_σ .*

References

- [1] Adjan, S. I., Novikov, P. S.: On a semicontinuous function. Moskov. Gos. Ped. Inst. Uchen. Zap. 138, 3–10, (1958) (in Russian)
- [2] Banach, T., Bokalo, B.: On scatteredly continuous maps between topological spaces. Topology and its Applications 157, 108–122 (2010)
- [3] Davies, R. O., Tricot, C.: A theorem about countable decomposability. Mathematical Proceedings of the Cambridge Philosophical Society, (03) **91**, 457–458 (1982)
- [4] Hansell, R. W., Jayne, J. E., Rogers, C. A.: Piece-wise closed functions and almost discretely σ -decomposable families. Mathematika 32, 229–247 (1985)
- [5] Holický, P., Spurný, J.: F_σ -mappings and the invariance of absolute Borel classes. Fund. Math. 182, 193–204 (2004)
- [6] Jayne, J. E., Rogers, C. A.: Piece-Wise Closed Functions. Math. Ann. 255, 499–518 (1981)
- [7] Jayne, J. E., Rogers, C. A.: Piece-Wise Closed Functions: Corrigendum. Math. Ann. 267, 143 (1984)
- [8] Keldiš, L.: Sur les fonctions premières mesurables B . Dokl. Akad. Nauk. SSSR, 4, 192–197 (1934)
- [9] Ostrovsky, A.: An alternative approach to the decomposition of functions. Topology and its Applications, 159, 200–208 (2012)
- [10] Ostrovsky, A.: The structure of LC -continuous functions. Acta Mathematica Hungarica, 133, 372–375 (2011)

UNIBW, MUNICH, GERMANY

E-mail address: Alexey.Ostrovskiy@UniBw.de